

# WOLF TRAX 57.5 % COPPER DDP™

Guaranteed Analysis:

Copper (Cu) ..... 57.5%

Dry Dispersible Powder formulated from copper sulfate, copper oxide and other inert ingredients.

Contains 244.4 lbs copper per 425 lb package. Contains 111 kg copper per 193 kg package.

## For the correction of copper deficiencies in crops

KEEP OUT OF REACH OF CHILDREN. WEAR GOGGLES, GLOVES AND DUST MASK WHEN HANDLING. Use only outdoors or in a well-ventilated area. Do not eat, drink or smoke when using this product. Avoid breathing dust or prolonged contact with skin or eyes. Wash thoroughly after handling. Safety data sheet available. If swallowed: Call a poison center/doctor if you feel unwell. Rinse mouth. If on skin: Wash with plenty of water. Take off immediately all contaminated clothing and wash it before reuse. If skin irritation occurs: Get medical advice/attention. If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention. If inhaled: Remove person to fresh air and keep comfortable for breathing. Call a poison center/doctor if you feel unwell. This product is a "Hazardous Chemical" as defined by the OSHA Hazard Communication Standard, 29 CFR 1910.1200. PRECAUTIONS. Always read and follow labelled directions for application. Over application can result in phytotoxicity to crops. STORAGE AND DISPOSAL. Store in a tightly closed container, in a cool dry well-ventilated facility with no access to the general public. Do not store near feed, foodstuffs or in proximity to water bodies. If spills occur, clean up by sweeping up. Dispose of wastes or containers at an approved waste disposal facility. Dispose of containers by thoroughly emptying powder into tank and disposing of packaging in a sanitary landfill or at a waste disposal facility as recommended by state or local authorities. WARRANTY DISCLAIMER. The seller shall not be liable for special, incidental or consequential damages or for damages in the form of penalties. The seller makes no other claims to the buyer for damages arising out of misuse of the product or the failure to follow label instructions. The goods sold to you are subject only to the guarantee specified on the label. The seller makes no other claims to the buyer either expressed or implied regarding the product sold with respect to the fitness of the product for a particular purpose. Ineffectiveness or unintended consequences may result as a consequence of manner of use of the product, which is beyond the control of the seller. In no case will the seller be held liable for damages arising as a result of mishandling of the product. The buyer assumes all such risks. The user prior to application must test compatibility of the product with other chemicals and the seller does not warrant that the product is compatible with all chemicals under all conditions. Information regarding the contents and levels of metals in this product is available on the Internet at <http://regulatory-info-wt.com>.

## DIRECTIONS FOR USE

**Coating on Fertilizer Blends** - This product may be used as a coating on all fertilizer blends (including urea or urea blends). Application rates and timings should be based on a soil and/or tissue test, as well as consultations with a professional agricultural consultant. Apply up to 1 lb of Wolf Trax Nutrient per 100 lbs of fertilizer (1 kg per 100 kg of fertilizer). Combined amounts of all Wolf Trax Nutrients added beyond 1 lb per 100 lbs of fertilizer may not adhere to the fertilizer. To apply, pour the required rate on to the fertilizer as it is being blended. Allow the product to mix thoroughly with the blend for several minutes.

**Mixing instructions for foliar applications** - Fill tank to 1/3 full, start agitation and pour package contents into tank. Use sufficient water to ensure thorough coverage. If mixing with a crop protection product, always add the micronutrient to the spray solution before adding the crop protection product. Prior to spraying, conduct a compatibility test by adding the products to water in a quart jar to ensure that the products will remain in solution. It is also recommended to use 50 mesh or coarser screens on all nozzles. **Foliar application rates and timing** -The following rates are suggestions based on correcting deficiencies seen in extensive field-testing. Individual rate recommendations may be obtained using a tissue test.

Crop	Foliar Rate	Foliar Timing	 <p>Warning H302: Harmful if swallowed. H315: Causes skin irritation. H319: Causes serious eye irritation.</p> <p>94% of the mixture consists of component(s) of unknown acute inhalation toxicity. 4.4% of the mixture consists of component(s) of unknown acute oral toxicity.</p>
Apples, pears, cherries, plums, citrus, almonds, nuts, peaches, avocado, figs	11 oz/acre (770 grams/ha)	Apply to foliage as a cover spray in spring. Repeat application if necessary up until bloom. (May also be applied to soil in liquid fertilizer at rates of up to 2.5 lbs/acre (2.8 kg/ha))	
Grapes, strawberries, cane berries, melons	11 oz/acre (770 grams/ha)	Apply before bloom. If an additional application is necessary, use 5.5 oz/acre (390 grams/ha).	
Potatoes	4 oz/acre (300 grams/ha)	Apply in early spring. Repeat application within four weeks if necessary.	
Vegetable crops	5.5 oz/acre (390 grams/ha)	Apply in early spring. Repeat application if necessary. Apply before blooming.	
Cereals, canola, flax	4 oz/acre (300 grams/ha)	Apply within 30 days of seeding for best results. May be applied up until the flag leaf has emerged (cereals) or flowering.	
Corn, cotton, sorghum	4 oz/acre (300 grams/ha)	Apply in early spring. Repeat application if necessary. Apply before blooming.	
Peanuts	4 oz/acre (300 grams/ha)	Apply just prior to bloom.	
Beans, soybeans	4 oz/acre (300 grams/ha)	Apply in early spring. For best results apply by the 4-trifoliolate leaf stage. Repeat application if necessary. Apply before blooming.	
Alfalfa, turfgrass	11 oz/acre (770 grams/ha)	Apply from three weeks after growth has resumed. Repeat application if necessary.	
Rice	11 oz/acre (770 grams/ha)	Apply using enough water for thorough coverage. Apply prior to pollination.	

**Manufactured For:** Compass Minerals USA Inc. 9900 W. 109th Street, Suite 100 Overland Park, KS 66210 Toll free 855.237.9653 [www.wolftrax.com](http://www.wolftrax.com)