

WOLF TRAX 47% IRON DDP™

Guaranteed Analysis

Iron (Fe) 47%

Dry Dispersible Powder formulated from ferrous sulfate, red iron oxide and other inert ingredients.

Contains 159.8 lbs iron per 340 lb package. Contains 72.4 kg iron per 154 kg package.

For the correction of iron deficiencies in crops. KEEP OUT OF REACH OF CHILDREN. WEAR GOGGLES, GLOVES AND DUST MASK WHEN HANDLING. Use only outdoors or in a well-ventilated area. Do not eat, drink or smoke when using this product. Avoid breathing dust or prolonged contact with skin or eyes. Wash thoroughly after handling. Safety data sheet available. If swallowed: Call a poison center/doctor if you feel unwell. Rinse mouth. If on skin: Wash with plenty of water. Take off immediately all contaminated clothing and wash it before reuse. If skin irritation occurs: Get medical advice/attention. If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention. If inhaled: Remove person to fresh air and keep comfortable for breathing. Call a poison center/doctor if you feel unwell. PRECAUTIONS. Always read and follow labelled directions for application. Over application can result in phytotoxicity to crops. STORAGE AND DISPOSAL. Store in a tightly closed container, in a cool dry well-ventilated facility with no access to the general public. Do not store near feed, foodstuffs or in proximity to water bodies. If spills occur, clean up by sweeping up. Dispose of wastes or containers at an approved waste disposal facility. Dispose of containers by thoroughly emptying powder into tank and disposing of packaging in a sanitary landfill or at a waste disposal facility as recommended by state or local authorities. This product is a "Hazardous Chemical" as defined by the OSHA Hazard Communication Standard, 29 CFR 1910.1200. Information regarding the contents and levels of metals in this product is available on the Internet at <http://regulatory-info-wt.com>. WARRANTY DISCLAIMER. The seller shall not be liable for special, incidental or consequential damages or for damages in the form of penalties. The seller makes no other claims to the buyer for damages arising out of misuse of the product or the failure to follow label instructions. The goods sold to you are subject only to the guarantee specified on the label. The seller makes no other claims to the buyer either expressed or implied regarding the product sold with respect to the fitness of the product for a particular purpose. Ineffectiveness or unintended consequences may result as a consequence of manner of use of the product, which is beyond the control of the seller. In no case will the seller be held liable for damages arising as a result of mishandling of the product. The buyer assumes all such risks. The user prior to application must test compatibility of the product with other chemicals and the seller does not warrant that the product is compatible with all chemicals under all conditions.

DIRECTIONS FOR USE

Coating on Fertilizer Blends - This product may be used as a coating on all fertilizer blends (including urea or urea blends). Application rates and timings should be based on a soil and/or tissue test, as well as consultations with a professional agricultural consultant. Apply up to 1 lb of Wolf Trax Nutrient per 100 lbs of fertilizer (1 kg per 100 kg of fertilizer). Combined amounts of all Wolf Trax Nutrients added beyond 1 lb per 100 lbs of fertilizer may not adhere to the fertilizer. To apply, pour the required rate on to the fertilizer as it is being blended. Allow the product to mix thoroughly with the blend for several minutes.

Mixing instructions for foliar applications - Fill tank to 1/3 full, start agitation and pour required amount into tank. Use sufficient water to ensure thorough coverage. If mixing with a crop protection product, always add the micronutrient to the spray solution before adding the crop protection product. Prior to spraying, conduct a compatibility test by adding the products to water in a quart jar to ensure that the products will remain in solution. It is also recommended to use 50 mesh or coarser screens on all nozzles. **Foliar application rates and timing** – The following rates are based on correcting deficiencies seen in extensive field-testing. Individual rate recommendations may be obtained using a tissue test. Apply when the foliage is actively growing. For best results, apply after irrigation or rainfall.

| Crop | Rates and timing |
|--|---|
| Fruit and nut crops such as apples, cherries, pears, plums, citrus, almonds | Soil applications – Apply 1.2 lbs/ acre (1,350 grams/ha) coated onto granular fertilizer. Foliar applications – Apply 13 oz/acre (910 grams/ha) in sufficient water to thoroughly cover leaves. If necessary, repeat at petal fall. Caution – Use a maximum of 2.4 lbs/acre (2.7 kg/ha) per season in plums or sensitive varieties. |
| Vine and berry crops such as grapes, raspberries, strawberries | Soil applications - Apply 13 oz/acre (910 grams/ha) coated onto granular fertilizer Foliar applications – Apply up to 13 oz/acre (910 grams/ha) in sufficient water to thoroughly cover leaves before bloom. |
| Field crops such as beans, cereals, corn, cotton, peas, potatoes, sorghum, canola, flax | Apply 4 oz/acre (300 grams/ha) in soil or as a foliar spray. Repeat as necessary. For best results with foliar applications in field crops, apply as follows; (1) bean crops – between the 2 nd and 4 th trifoliolate leaf, (2) corn or cotton – within 3 weeks of emergence, (3) barley – within 30 days of planting, (4) potatoes – within 3 weeks of emergence |
| Vegetable crops | Apply 7 oz/acre (495 grams/ha) on soil or as a foliar spray within 3 weeks of emergence. Repeat as necessary. |
| Ornamentals such as potted shrubs, trees, and flowers as well as field grown nursery stock | Apply 1 oz per 1000 ft ² (24 grams per 100 m ²) as a foliar application. Repeat as necessary. |
| Turfgrass | Apply 4 oz per acre (300 grams/ha) as a foliar application. Repeat as necessary. |
| Rice | Apply 4 oz per acre (300 grams/ha) in sufficient water to ensure thorough coverage, and repeat as necessary. Apply prior to pollination |



Warning H315: Causes skin irritation. H319: Causes serious eye irritation. H335: May cause respiratory irritation

100% of the mixture consists of component(s) of unknown acute inhalation toxicity.
14% of the mixture consists of component(s) of unknown acute oral toxicity.

Manufactured For: Compass Minerals USA Inc. 9900 W. 109th Street, Suite 100 Overland Park, KS 66210 Toll free 855.237.9653 www.wolftrax.com

WTFEUS024_HF

Patent Issued USA Patent # 7,410,522
Patent Issued USA Patent # 7,445,657

Net Wt. – 340 lbs (154 kg)